

BIELLIPTIC CURVES OF GENUS 3 IN THE HYPERELLIPTIC MODULI

T. SHASKA AND F. THOMPSON

ABSTRACT. In this paper we study bielliptic curves of genus 3 defined over an algebraically closed field k and the intersection of the moduli space \mathcal{M}_3^h of such curves with the hyperelliptic moduli \mathcal{H}_3 . Such intersection \mathcal{S} is an irreducible, 3-dimensional, rational algebraic variety. We determine the equation of this space in terms of the $GL(2, k)$ -invariants of binary octavics as defined in [27] and find a birational parametrization of \mathcal{S} . We also compute all possible subloci of curves for all possible automorphism group G . Moreover, for every rational moduli point $\mathfrak{p} \in \mathcal{S}$, such that $|\text{Aut}(\mathfrak{p})| > 4$, we give explicitly a rational model of the corresponding curve over its field of moduli in terms of the $GL(2, k)$ -invariants. genus 3 hyperelliptic curves and dihedral invariants and absolute invariants

1. INTRODUCTION

The moduli space \mathcal{M}_g of algebraic curves of genus $g \geq 2$, defined over an algebraically closed field k , is an interesting object that has received plenty of attention since the mid XX-century. It is an irreducible quasi-projective variety of dimension $3g - 3$. Understanding the stratification of this space has been also a major problem with many papers written on the subject to this day. There are two main difficulties on this problem:

- i) an explicit description of \mathcal{M}_g is not known (i.e., a coordinate in \mathcal{M}_g),*
- ii) a list of automorphism groups for a fixed $g \geq 2$ has not been known.*

Naturally one has a better chance to address the above problem if focused on the hyperelliptic sublocus \mathcal{H}_g of \mathcal{M}_g , since it is easier to pick a coordinate on the space \mathcal{H}_g . After all, the hyperelliptic curves were well understood since the XIX-century and restricting the problem to the hyperelliptic locus seems reasonable. It was well known to classical algebraic geometers of the XIX-century that the isomorphism classes of hyperelliptic curves defined over an algebraically closed field k correspond to the orbits of the $GL_2(k)$ action on the space of binary forms of degree $2g + 2$ with coefficients from k . This was, among others, one of the main motivations of the invariant theory during the XIX-century. For the generalization to the case of superelliptic curves one can check [21].

The case of genus 2 had been studied extensively by XIX-century mathematicians; see [6, 7] even though the concept of the moduli space was not quite refined at the time. About a decade ago Gaudry/Schost in [13] attacked the problem for $g = 2$ from the computational point of view. After all, a coordinate in \mathcal{M}_2 could be fixed using the Igusa invariants and the list of automorphism groups of genus 2 was known; see [14] among others. At the same time that [13] was being circulated as a preprint, Shaska/Völklein [30] considered the problem from a more

group-theoretical point of view. Of course, the main case in both those papers was the case when the genus 2 curves had an elliptic involution. The locus \mathcal{L}_2 of such curves is a 2-dimensional irreducible variety in \mathcal{M}_2 computed in both papers. In the process, a group action was discovered in [30] and its u, v invariants were instrumental in computing equations for the strata of \mathcal{M}_2 . The map

$$(1) \quad \phi_2(u, v) \rightarrow (i_1, i_2, i_3)$$

provides a birational parametrization of the space \mathcal{L}_2 , where i_1, i_2, i_3 are $GL_2(k)$ -invariants in the space of binary sextics. The singular locus of this map correspond to genus 2 curves with larger automorphism group; see [30, Lemma 3]. The paper [30] spurred interest in two directions. First, it naturally brought to the attention of the authors the problem of automorphism groups of curves of genus $g \geq 2$. This corresponds to the ii) part of the problem stated in the beginning. Second, naturally raised the question whether the invariants u, v for $g = 2$ could be generalized to higher genus. In the next two paragraphs we consider each direction in more details.

Determining the list of automorphism groups of algebraic curves of genus $g \geq 2$ is a classical problem. There were hundreds of papers in the subject before 2001, most of them considering specific cases for small genus. However, there was one interesting development at the time that it seems as it did not get the attention it deserved. Breuer computed all signatures of the groups acting on compact Riemann surfaces for genus $g \leq 48$; see [5]. The restriction $g \leq 48$ is merely technical and Breuer's algorithm works for any genus, providing that some careful analysis is required for sporadic cases. Using results in [5] and the theory of Hurwitz spaces, Magaard, Shpectorov, Shaska, Völklein determined an algorithm of how to determine the list of full automorphism groups of curves for any given genus $g \geq 2$; see [20]. In [20] a complete list of full automorphism groups for curves of genus 3 was determined and the corresponding equations were provided as a way to illustrate the methods described in that paper. There were tens of papers on the case of genus $g = 3$ before [20] appeared. Moreover, by methods in [20] the list of full automorphism groups of curves for any genus $g \geq 2$ can be determined. This settles the second part ii) of the initial problem.

The second direction that was spurred by [30] was the problem of generalizing the map (1) to higher genus. Natural questions to follow would be whether the curves with larger automorphism groups would be in the singular locus of ϕ . The group action discovered in [30] was generalized in [26] and then in a more formal paper in [16] where such invariants in higher genus were called **dihedral invariants**. For a genus $g \geq 2$ now we have a map

$$(2) \quad \phi_g(\mathfrak{s}_4, \dots, \mathfrak{s}_g) \rightarrow (t_1, \dots, t_{2g-1}),$$

where t_1, \dots, t_{2g-1} are $GL_2(k)$ -invariants in the space of binary forms of degree $2g - 1$. In papers [24], [15], and [16] the case of stratification of the hyperelliptic moduli \mathcal{H}_3 was treated to illustrate the general theory. Dihedral invariants have been used quite extensively since by many authors. They were generalized for fields of positive characteristic by [2] and are defined again in the projective version in [19], where they are renamed as *dihedral arithmetic invariants*.

This paper takes another look at the study of genus 3 hyperelliptic curves with extra automorphisms. The general strategy for $g = 3$ was quite obvious a decade ago; one computes the Shioda invariants defined in [31] starting from the Table 1 of [20] and then eliminating the parameters which appear as coefficients of the

curve. Such computations were simplified considerable via the dihedral invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$. There was an obvious drawback compared to the genus $g = 2$ case; the $GL_2(k)$ -invariants for $g = 3$ were not known. Hence, the obvious strategy was to describe the strata in terms of the $SL_2(k)$ -invariants defined by Shioda in [31]. This approach is taken in [18]. It is not clear from [18] if the dihedral invariants were used in these computations or they were performed straight from the equations of the curves as in Table 3 in [20]. In any case, Shioda invariants J_2, \dots, J_{10} are computed in [18] and using syzygies determined in [31, Theorem 5] the authors determine each loci in \mathcal{H}_3 . In [27] it was shown that the syzygies determined by Shioda in [31, Theorem 5] are not correct. It is unclear if the authors in [18] have corrected such syzygies, otherwise all the results of [18] could be incorrect.

The motivation for this paper was the definition of $GL_2(k)$ -invariants in [27] where an explicit equation of the hyperelliptic moduli \mathcal{H}_3 is given in the ambient space \mathbb{C}^6 . Using the absolute invariants t_1, \dots, t_6 as in [27] and the dihedral invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ one can easily compute the locus in \mathcal{H}_3 for each case of Table 3 of [20]. The drawback of invariants t_1, \dots, t_6 is that they are not defined everywhere. However, this is done by choice so that their degrees are kept small. This makes computations a lot easier. One can get projective equations (i.e., equations in terms of J_2, \dots, J_8) from our equations simply by replacing t_1, \dots, t_6 with their definitions and clearing denominators.

The paper is organized as follows. In section 2 we give a brief description of the invariants of the binary octavics and definitions of absolute invariants. Notice that our definitions of J_2, \dots, J_8 are slightly different from those of Shioda. In section 3 we discuss genus 3 hyperelliptic curves with an elliptic involution and derive an parametric equation for such family of curves. This is rather known material that has been treated in [16, 26].

Section 4 is the main section of the paper where the equation for the locus of curves with an elliptic involution. The main theorem here describes the equation of the irreducible sub variety \mathcal{S} in the hyperelliptic moduli \mathcal{H}_3 . We show that for a generic curve C in \mathcal{S} the field of definition is at most a degree 2 extension of the field of moduli. This is an improvement from [18] where it is shown that this bound is eight. For example, for the case when the group is V_4 , we get a model for the parametric curve defined over a quadratic extension of the field of moduli versus a degree 8 extension in [18].

In section 5, we determine the equation of all 1 and 2-dimensional loci for any fixed automorphism group G . Parametrization of such loci were also given in [15]. Here we compute them in terms of the absolute invariants t_1, \dots, t_6 .

The goal of this paper was to describe the stratification of the space \mathcal{H}_3 in terms of the absolute invariants t_1, \dots, t_6 . The benefit of this approach is that there are fewer equations and even simpler ones. The results in [27] make it possible that we do not have to use the invariants J_9, J_{10} and have fewer equations in each case. We get better results compared to [18] in the case of the group V_4 and \mathbb{Z}_2^3 on the minimal equation of the curves over their field of moduli.

In the case of group \mathbb{Z}_2^3 we prove that the field of moduli is a field of definition and give a model of the curve over its field of definition. Some of these results were not new to us, since they were proved in [16]. However, in this paper we are able to explicitly describe such results in terms of the absolute invariants t_1, \dots, t_6 . All our results are implemented in a Maple package which is provided for free on [29].

Notation: Throughout this paper, by a "curve" we mean an irreducible algebraic curve defined over an algebraically closed field k . While we use invariants J_2, \dots, J_8 of binary octavics as Shioda [31], the reader must be aware that our definitions are not the same as those in Shioda's paper, instead we use the definitions as [27]. We also use the dihedral invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ which are the same as those used in [15] u, v, w , where $\mathfrak{s}_4 = u$, $\mathfrak{s}_3 = v$, $\mathfrak{s}_2 = w$.

2. BIELLIPTIC GENUS 3 CURVES

Let k be an algebraically closed field of characteristic zero and \mathcal{X} an irreducible, smooth, projective curve of genus $g \geq 3$ defined over k . As usual, we denote by \mathcal{M}_g the coarse moduli space of smooth curves of genus $g \geq 2$ and by \mathcal{H}_g the hyperelliptic locus in \mathcal{M}_g . The isomorphism class of C , i.e. the corresponding point in \mathcal{M}_g , is denoted by $[C]$.

A curve C is called *bielliptic* if it admits a degree 2 morphism $\pi : C \rightarrow E$ onto an elliptic curve. Let

$$\mathcal{M}_g^b = \{[C] \in \mathcal{M}_g : C \text{ bielliptic} \}$$

be the locus of bielliptic curves in \mathcal{M}_g . \mathcal{M}_g^b is an irreducible $(2g - 2)$ -dimensional sub variety of \mathcal{M}_g . For $g = 3$ \mathcal{M}_3^b is the unique component of maximal dimension of the singular locus $Sing(\mathcal{M}_3)$; see [8, 9] for details. It is known that

i) \mathcal{M}_3^b is rational

ii) $\mathcal{M}_3^b \cap \mathcal{H}_3$ is an irreducible, codimension 1, rational subvariety of \mathcal{M}_3^b ,

see [3, Theorem 1.1] for details. In this paper we aim to find an algebraic equation for the space $\mathcal{S} := \mathcal{M}_3^b \cap \mathcal{H}_3$. An algebraic equation for \mathcal{M}_3^b using invariants of ternary quartics and a theorem of Kovalevskaia is intended in [11].

Let α be the element in $\text{Aut}(C)$ which interchanges the sheets of $\pi : C \rightarrow E$ such that $E \cong C/\langle \alpha \rangle$. We call α the **elliptic involution** of C corresponding to π . Hence, the space $\mathcal{S} = \mathcal{M}_3^b \cap \mathcal{H}_3$ is exactly the space of genus 3 hyperelliptic curves with elliptic involutions. Such space has been studied before from the point of view of automorphism groups, as described in details in the introduction.

For a fixed group G acting on a genus g algebraic curves \mathcal{X}_g we have a covering $\mathcal{X}_g \rightarrow \mathcal{X}_g/G$. All possible ramification structures of such covering for any genus g hyperelliptic curves were determined in [22]. Indeed, this is also done for all superelliptic curves; see [4] for details. In the case of hyperelliptic curves of genus 3, each group occurs only with one signature; see [20]. Hence, there is no confusion if we denote by $\mathcal{S}(G)$ the locus in \mathcal{H}_3 of all curves with automorphism group isomorphic to G (i.e., $G \hookrightarrow \text{Aut}(\mathcal{X}_g)$). The loci $\mathcal{S}(G)$ is not a priori irreducible. In general, irreducibility is checked by the braid action on Nielsen tuples. The locus $\mathcal{S}(G)$ is a Hurwitz space of covers with monodromy group G and fixed ramification structure. However, under our assumptions ($g = 3$ and hyperelliptic) this is always the case as shown in [22] and we will avoid that discussion here.

3. GENUS 3 HYPERELLIPTIC FIELDS WITH ELLIPTIC INVOLUTIONS

Let K be a genus 3 hyperelliptic field. Then K has exactly one genus 0 subfield of degree 2, call it $k(X)$. It is the fixed field of the **hyperelliptic involution** ω_0 in $\text{Aut}(K)$. Thus, ω_0 is central in $\text{Aut}(K)$, where $\text{Aut}(K)$ denotes the group $\text{Aut}(K/k)$. It induces a subgroup of $\text{Aut}(k(X))$ which is naturally isomorphic

to $\overline{\text{Aut}}(K) := \text{Aut}(K)/\langle \omega_0 \rangle$. The latter is called the **reduced automorphism group** of K .

If ω_1 is a non-hyperelliptic involution in G then $\omega_2 := \omega_0 \omega_1$ is another one. So the non-hyperelliptic involutions come naturally in (unordered) pairs ω_1, ω_2 . These pairs correspond bijectively to the Klein 4-groups in G . Indeed, each Klein 4-group in G contains ω_0 .

Definition 1. *We will consider pairs (K, ε) with K a genus 3 hyperelliptic field and ε an non-hyperelliptic involution in \bar{G} . Two such pairs (K, ε) and (K', ε') are called isomorphic if there is a k -isomorphism $\alpha : K \rightarrow K'$ with $\varepsilon' = \alpha \varepsilon \alpha^{-1}$.*

Let ε be an non-hyperelliptic involution in \bar{G} . We can choose the generator X of $\text{Fix}(\omega_0)$ such that $\varepsilon(X) = -X$. Then $K = k(X, Y)$ where X, Y satisfy equation

$$Y^2 = (X^2 - \alpha_1^2)(X^2 - \alpha_2^2)(X^2 - \alpha_3^2)(X^2 - \alpha_4^2)$$

for some $\alpha_i \in k$, $i = 1, \dots, 4$. Denote by

$$(3) \quad \begin{aligned} s_1 &= -(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) \\ s_2 &= (\alpha_1 \alpha_2)^2 + (\alpha_1 \alpha_3)^2 + (\alpha_1 \alpha_4)^2 + (\alpha_2 \alpha_3)^2 + (\alpha_2 \alpha_4)^2 + (\alpha_3 \alpha_4)^2 \\ s_3 &= -(\alpha_1 \alpha_2 \alpha_3)^2 - (\alpha_4 \alpha_1 \alpha_2)^2 - (\alpha_4 \alpha_3 \alpha_1)^2 - (\alpha_4 \alpha_3 \alpha_2)^2 \\ s_4 &= -(\alpha_1 \alpha_2 \alpha_3 \alpha_4)^2 \end{aligned}$$

Then, we have

$$Y^2 = X^8 + s_1 X^6 + s_2 X^4 + s_3 X^2 + s_4$$

with $s_1, s_2, s_3, s_4 \in k$, $s_4 \neq 0$. Further $E_1 = k(X^2, Y)$ and $C = k(X^2, YX)$ are the two subfields corresponding to ε of genus 1 and 2 respectively.

Preserving the condition $\varepsilon(X) = -X$ we can further modify X such that $s_4 = 1$. Then, we have the following:

Lemma 1. *Every genus 3 hyperelliptic curve \mathcal{X} , defined over a field k , which has an non-hyperelliptic involution has equation*

$$(4) \quad Y^2 = X^8 + aX^6 + bX^4 + cX^2 + 1$$

for some $a, b, c \in k^3$, where the polynomial on the right has non-zero discriminant.

Indeed, the non-hyperelliptic involution above is an elliptic involution and \mathcal{X} is bielliptic. There is another non-hyperelliptic involution of \mathcal{X} , as noted above, namely $\omega_2 := \omega_0 \omega_1$ which fixes a genus 2 field. See [28] for the equation of this genus 2 subfield and the arithmetic of such curves. hence, we have the following result; see [3] or [28] for details.

Proposition 1. *$[C] \in \mathcal{S}$ if and only if C is a double covering of a genus 2 curve.*

The above conditions determine X up to coordinate change by the group $\langle \tau_1, \tau_2 \rangle$ where

$$\tau_1 : X \rightarrow \zeta_8 X, \quad \text{and} \quad \tau_2 : X \rightarrow \frac{1}{X},$$

and ζ_8 is a primitive 8-th root of unity in k . Hence,

$$\tau_1 : (a, b, c) \rightarrow (\zeta_8^6 a, \zeta_8^4 b, \zeta^2 c),$$

and

$$\tau_2 : (a, b, c) \rightarrow (c, b, a).$$

Then, $|\tau_1| = 4$ and $|\tau_2| = 2$. The group generated by τ_1 and τ_2 is the dihedral group of order 8. Invariants of this action are

$$(5) \quad \begin{aligned} \mathfrak{s}_2 &= ac, \\ \mathfrak{s}_3 &= (a^2 + c^2)b, \\ \mathfrak{s}_4 &= a^4 + c^4, \end{aligned}$$

since

$$\begin{aligned} \tau_1(a^4 + c^4) &= (\zeta_8^6 a)^4 + (\zeta_8^2 c)^4 = a^4 + c^4 \\ \tau_1((a^2 + c^2)b) &= (\zeta_8^4 a^2 + \zeta_8^4 c^2) \cdot (\zeta_8^4 b) = (a^2 + c^2)b \\ \tau_1(ac) &= \zeta_8^6 a \cdot \zeta_8^2 c = ac \end{aligned}$$

Since they are symmetric in a and c , then they are obviously invariant under τ_2 . Notice that $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ are homogenous polynomials of degree 2, 3, and 4 respectively. The subscript i represents the degree of the polynomial \mathfrak{s}_i .

Since the above transformations are automorphisms of the projective line $\mathbb{P}^1(k)$ then the $SL_2(k)$ invariants must be expressed in terms of $\mathfrak{s}_4, \mathfrak{s}_3$, and \mathfrak{s}_2 . In these parameters, the discriminant of the octavic polynomial on the right hand side of Eq. (4) equals $-\frac{256}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^4} \Delta^2$, where

$$(6) \quad \begin{aligned} \Delta &= 132\mathfrak{s}_2^4 \mathfrak{s}_4 - 18\mathfrak{s}_4^2 \mathfrak{s}_2 \mathfrak{s}_3 - 72\mathfrak{s}_4 \mathfrak{s}_2^3 \mathfrak{s}_3 - \mathfrak{s}_4 \mathfrak{s}_2^2 \mathfrak{s}_3^2 + 80\mathfrak{s}_2 \mathfrak{s}_3^2 \mathfrak{s}_4 - 576\mathfrak{s}_3 \mathfrak{s}_2^2 \mathfrak{s}_4 \\ &\quad - 256\mathfrak{s}_4^2 + 768\mathfrak{s}_4 \mathfrak{s}_2^3 - 1024\mathfrak{s}_4 \mathfrak{s}_2^2 + 256\mathfrak{s}_2^2 \mathfrak{s}_3^2 - 576\mathfrak{s}_2^4 \mathfrak{s}_3 + 768\mathfrak{s}_2^5 + 24\mathfrak{s}_2^6 \\ &\quad - 16\mathfrak{s}_3^4 - 1024\mathfrak{s}_2^4 + 128\mathfrak{s}_3^2 \mathfrak{s}_4 + 192\mathfrak{s}_4^2 \mathfrak{s}_2 + 114\mathfrak{s}_4^2 \mathfrak{s}_2^2 + 4\mathfrak{s}_4^2 \mathfrak{s}_2^3 - 144\mathfrak{s}_4^2 \mathfrak{s}_3 \\ &\quad + 16\mathfrak{s}_4 \mathfrak{s}_2^5 - 72\mathfrak{s}_2^5 \mathfrak{s}_3 - 2\mathfrak{s}_2^4 \mathfrak{s}_3^2 + 160\mathfrak{s}_2^3 \mathfrak{s}_3^2 + 4\mathfrak{s}_3^3 \mathfrak{s}_4 + 8\mathfrak{s}_3^3 \mathfrak{s}_2^2 + 27\mathfrak{s}_4^3 + 16\mathfrak{s}_2^7 \end{aligned}$$

The map

$$(a, b, c) \mapsto (\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$$

is a branched Galois covering with group D_4 of the set

$$\{(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) \in k^3 : \Delta_{(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)} \neq 0\}$$

by the corresponding open subset of a, b, c -space. In any case, it is true that if a, b, c and a', b', c' have the same $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ -invariants then they are conjugate under $\langle \tau_1, \tau_2 \rangle$.

The case when $\mathfrak{s}_3 = 0$ must be treated separately. We have two sub cases $a^2 + c^2 = 0$ or $b = 0$. Then we define new invariants as follows:

$$(7) \quad \mathfrak{p}(\mathcal{X}_3) = \begin{cases} w = b^2 & \text{if } a = c = 0, \\ (\mathfrak{s}_2, w, \mathfrak{s}_4) & \text{if } a^2 + c^2 = 0 \text{ and } b \neq 0, \\ (\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) & \text{otherwise.} \end{cases}$$

The invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4, \dots$ are valid for any genus $g \geq 2$ and are called by many authors **dihedral invariants**. They were discovered by the second author in his PhD thesis and appeared for the first time in the literature in Shaska/Völklein [30]. Then, they appeared for genus $g = 3$ in [15, 24] and were generalized for every genus in [16]. They were generalized a ditto to all cyclic curves by Antoniadis/Kontogiorgis [2]. In [19] a projective version of these dihedral invariants are called *dihedral arithmetic invariants*.

Lemma 2. For $(a, b, c) \in k^3$ with $\Delta \neq 0$, equation (4) defines a genus 3 hyperelliptic field $K_{a,b,c} = k(X, Y)$. Its reduced automorphism group contains the elliptic involution $\varepsilon_{a,b,c} : X \mapsto -X$. Two such pairs $(K_{a,b,c}, \varepsilon_{a,b,c})$ and $(K_{a',b',c'}, \varepsilon_{a',b',c'})$ are isomorphic if and only if

$$(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) = (\mathfrak{s}'_2, \mathfrak{s}'_3, \mathfrak{s}'_4)$$

where $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ and $\mathfrak{s}'_2, \mathfrak{s}'_3, \mathfrak{s}'_4$ are dihedral invariants associated with a, b, c and a', b', c' , respectively.

Proof. An isomorphism α between these two pairs yields $K = k(X, Y) = k(X', Y')$ with $k(X) = k(X')$ such that X, Y satisfy (4) and X', Y' satisfy the corresponding equation with a, b, c replaced by a', b', c' . Further, $\varepsilon_{a,b,c}(X') = -X'$. Thus X' is conjugate to X under $\langle \tau_1, \tau_2 \rangle$ by the above remarks. This proves the condition is necessary. It is clearly sufficient. \square

Remark 1. If $(2\mathfrak{s}_4 + \mathfrak{s}_2^2) = 0$, then this implies that $a = c = 0$. In this case the equation of the curve becomes

$$Y^2 = X^8 + bX^4 + 1,$$

which corresponds to the curves with automorphism group $\mathbb{Z}_2 \times D_8$, (cf. Eq. (27))

We have the following theorem

Theorem 1. Let $(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) \in k^3 \setminus \{\Delta = 0\}$. Then the following hold:

i) The ordered triples $(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$ bijectively parameterize the isomorphism classes of pairs (K, ε) where K is a genus 3 hyperelliptic field and ε an elliptic involution of $\text{Aut}(K)$. The j -invariant of the elliptic subfield of K associated with ε is given by

$$(8) \quad j = \frac{64}{M} \cdot \frac{(-4\mathfrak{s}_3^2 - 48\mathfrak{s}_4 - 24\mathfrak{s}_2^2 + 3\mathfrak{s}_3^3 + 6\mathfrak{s}_4\mathfrak{s}_2)^3}{(2\mathfrak{s}_4 + \mathfrak{s}_2^2)},$$

where

$$\begin{aligned} M = & 66\mathfrak{s}_4\mathfrak{s}_2^4 - 2048\mathfrak{s}_4^2 - 512\mathfrak{s}_2^4 - 2048\mathfrak{s}_4\mathfrak{s}_2^2 - 128\mathfrak{s}_3^4 + 1024\mathfrak{s}_3^2\mathfrak{s}_4 + 512\mathfrak{s}_3^2\mathfrak{s}_2^2 + 228\mathfrak{s}_4^2\mathfrak{s}_2^2 \\ & + 768\mathfrak{s}_4^2\mathfrak{s}_2 + 216\mathfrak{s}_4^3 + \mathfrak{s}_2^7 + 3\mathfrak{s}_2^6 + 4\mathfrak{s}_2^3\mathfrak{s}_4^2 + 4\mathfrak{s}_2^5\mathfrak{s}_4 - \mathfrak{s}_3^2\mathfrak{s}_4^2 + 768\mathfrak{s}_4\mathfrak{s}_2^3 + 160\mathfrak{s}_3^2\mathfrak{s}_2^3 \\ & + 192\mathfrak{s}_2^5 - 2\mathfrak{s}_3^2\mathfrak{s}_4\mathfrak{s}_2^2 + 320\mathfrak{s}_3^2\mathfrak{s}_4\mathfrak{s}_2 - 72\mathfrak{s}_4^2\mathfrak{s}_3\mathfrak{s}_2 - 72\mathfrak{s}_4\mathfrak{s}_2^3\mathfrak{s}_3 - 1152\mathfrak{s}_4\mathfrak{s}_2^2\mathfrak{s}_3 + 32\mathfrak{s}_3^3\mathfrak{s}_4 \\ & - 1152\mathfrak{s}_4^2\mathfrak{s}_3 - 288\mathfrak{s}_2^4\mathfrak{s}_3 + 16\mathfrak{s}_3^3\mathfrak{s}_2^2 - 18\mathfrak{s}_2^5\mathfrak{s}_3 \end{aligned}$$

ii) There is another involution $\omega_0\varepsilon \in \text{Aut}(K)$ which fixes a genus 2 curve \mathcal{X}_2 with equation

$$Y^2 = X(X^4 + aX^3 + bX^2 + cX + 1).$$

The isomorphism class of \mathcal{X}_2 is determined uniquely by the triple $(\mathfrak{s}_2, \mathfrak{s}_4, \mathfrak{s}_4)$ as in Eq. (9).

iii) The triples $(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$ parametrize the isomorphism classes of genus 3 hyperelliptic fields with $V_4 \hookrightarrow \text{Aut}(K)$.

Proof. i) The automorphism $\varepsilon \in \text{Aut}(K)$ fixes a degree 2 elliptic subfield E which has equation

$$Y^2 = x^4 + ax^3 + bx^2 + cx + 1$$

and j -invariant given in terms of a , b , and c . Using substitutions in Eq. (14) we get $j(E)$ as in Eq.(8).

ii) The quotient $\mathcal{X}/\langle\omega_0\varepsilon\rangle$ has genus 2. This follows straight from the Riemann-Hurwitz formula. The equation of this genus 2 curve is as claimed; see [28]. The isomorphism class of this genus 2 curve is determined by the absolute invariants (i_1, i_2, i_3) . In terms of the $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ they have the following expressions

(9)

$$i_1 = \frac{9(2\mathfrak{s}_1 + \mathfrak{s}_3^2)}{D^2} (\mathfrak{s}_3^4 - 80\mathfrak{s}_3^2 - 72\mathfrak{s}_2^2 - 2\mathfrak{s}_3^2\mathfrak{s}_1 - 24\mathfrak{s}_1\mathfrak{s}_2 - 12\mathfrak{s}_3^2\mathfrak{s}_2 + 2\mathfrak{s}_3^3 - 160\mathfrak{s}_1 + 4\mathfrak{s}_3\mathfrak{s}_1)$$

$$i_2 = \frac{27(2\mathfrak{s}_1 + \mathfrak{s}_3^2)^2}{D^3} (2\mathfrak{s}_3^3\mathfrak{s}_1 - 1116\mathfrak{s}_1\mathfrak{s}_2 + \mathfrak{s}_3^5 - 2240\mathfrak{s}_3^2 + 162\mathfrak{s}_2^2\mathfrak{s}_3 + 864\mathfrak{s}_2^2 + 216\mathfrak{s}_1^2 + 114\mathfrak{s}_3^2\mathfrak{s}_1 + 3\mathfrak{s}_3^4 - 558\mathfrak{s}_3^2\mathfrak{s}_2 - 4480\mathfrak{s}_1 + 624\mathfrak{s}_3^3 - 18\mathfrak{s}_3^3\mathfrak{s}_2 - 36\mathfrak{s}_1\mathfrak{s}_2\mathfrak{s}_3 + 1248\mathfrak{s}_3\mathfrak{s}_1)$$

$$i_3 = \frac{243}{1024} \frac{(2\mathfrak{s}_1 + \mathfrak{s}_3^2)^3}{D^5} (\mathfrak{s}_3^7 - 128\mathfrak{s}_2^4 - 2048\mathfrak{s}_1^2 + 768\mathfrak{s}_3^3\mathfrak{s}_1 - 2048\mathfrak{s}_3^2\mathfrak{s}_1 + 192\mathfrak{s}_3^5 - 512\mathfrak{s}_3^4 + 216\mathfrak{s}_1^3 + 3\mathfrak{s}_3^6 - 72\mathfrak{s}_1^2\mathfrak{s}_3\mathfrak{s}_2 + 320\mathfrak{s}_1\mathfrak{s}_2^2\mathfrak{s}_3 - 72\mathfrak{s}_1\mathfrak{s}_3^3\mathfrak{s}_2 - 1152\mathfrak{s}_1\mathfrak{s}_3^2\mathfrak{s}_2 - 2\mathfrak{s}_2^2\mathfrak{s}_1\mathfrak{s}_3^2 - 1152\mathfrak{s}_1^2\mathfrak{s}_2 + 1024\mathfrak{s}_1\mathfrak{s}_2^2 + 160\mathfrak{s}_2^2\mathfrak{s}_3^3 + 512\mathfrak{s}_2^2\mathfrak{s}_3^2 - 18\mathfrak{s}_3^5\mathfrak{s}_2 - 288\mathfrak{s}_3^4\mathfrak{s}_2 + 768\mathfrak{s}_3\mathfrak{s}_1^2 + 4\mathfrak{s}_3^3\mathfrak{s}_1^2 + 4\mathfrak{s}_3^5\mathfrak{s}_1 + 228\mathfrak{s}_3^2\mathfrak{s}_1^2 + 66\mathfrak{s}_3^4\mathfrak{s}_1 + 16\mathfrak{s}_2^3\mathfrak{s}_3^2 + 32\mathfrak{s}_2^3\mathfrak{s}_1 - \mathfrak{s}_2^2\mathfrak{s}_3^4),$$

where $D = -20\mathfrak{s}_1 - 10\mathfrak{s}_3^2 + 2\mathfrak{s}_3^3 + 4\mathfrak{s}_3\mathfrak{s}_1 - 3\mathfrak{s}_2^2$.

iii) This is a straight consequence of the first two parts. The cases when $|\text{Aut}(K)| > 4$ are treated in Thm. 4. \square

4. THE LOCUS \mathcal{S} OF GENUS 3 HYPERELLIPTIC CURVES WITH ELLIPTIC INVOLUTIONS

In this section, first we briefly define the invariants of binary octavics. We will use interchangeably the terms genus 3 hyperelliptic curve and genus 3 hyperelliptic field. There is a one to one equivalence between the isomorphic classes of genus 3 hyperelliptic curves and projective classes of equivalence of binary octavics. Thus, we have to describe some basic properties of binary octavics. The following material can be found on works of classical algebraic geometers; see Alagna [1], van Gall [32], et al or for a modern version one can check [10].

The ring of invariants of binary octavics was also studied by Shioda [31]. However, the syzygies among such $SL_2(k)$ -invariants described in the Shioda's paper seem to be incorrect. In [27] such $SL_2(k)$ -invariants J_2, \dots, J_8 are redefined and the algebraic relations among them determined. Furthermore, $GL_2(k)$ -invariants t_1, \dots, t_6 are defined and relation among them determined. Throughout this paper we will make use of these $GL_2(k)$ -invariants and therefore follow definitions from [27].

Let $f(X, Y)$ be the binary octavic

$$f(X, Y) = \sum_{i=0}^8 a_i X^i Y^{8-i}.$$

defined over an algebraically closed field k . We define the following covariants:

$$(10) \quad \begin{aligned} g &= (f, f)^4, & k &= (f, f)^6, & h &= (k, k)^2, \\ m &= (f, k)^4, & n &= (f, h)^4, & p &= (g, k)^4, & q &= (g, h)^4, \end{aligned}$$

where the operator $(\cdot, \cdot)^n$ denotes the n -th transvection of two binary forms; see [27] among many other references.

Then, the following

$$(11) \quad \begin{aligned} J_2 &= 2^2 \cdot 5 \cdot 7 \cdot (f, f)^8, & J_3 &= \frac{1}{3} \cdot 2^4 \cdot 5^2 \cdot 7^3 \cdot (f, g)^8, \\ J_4 &= 2^9 \cdot 3 \cdot 7^4 \cdot (k, k)^4, & J_5 &= 2^9 \cdot 5 \cdot 7^5 \cdot (m, k)^4, \\ J_6 &= 2^{14} \cdot 3^2 \cdot 7^6 \cdot (k, h)^4, & J_7 &= 2^{14} \cdot 3 \cdot 5 \cdot 7^7 \cdot (m, h)^4, \\ J_8 &= 2^{17} \cdot 3 \cdot 5^2 \cdot 7^9 \cdot (p, h)^4, & J_9 &= 2^{19} \cdot 3^2 \cdot 5 \cdot 7^9 \cdot (n, h)^4, \\ J_{10} &= 2^{22} \cdot 3^2 \cdot 5^2 \cdot 7^{11} (q, h)^4 \end{aligned}$$

are $SL_2(k)$ -invariants; see [27] for details. The following is a classical fact of invariant theory of binary forms.

Lemma 3. *Two binary forms $f(X, Y)$ and $f'(X, Y)$ are projectively equivalent via a matrix $M \in GL_2(k)$ if and only if*

$$J_i(f) = \lambda^i J_i(f'), \quad \text{where } \lambda = (\det(M))^4$$

The following technical result is helpful for the rest of the paper; see [27, Lemma 4].

Lemma 4. *If $J_2 = \dots = J_7 = 0$, then the binary octavic has a double root.*

Next, we define $GL(2, k)$ -invariants as follows

$$t_1 := \frac{J_3^2}{J_2^3}, \quad t_2 := \frac{J_4}{J_2^2}, \quad t_3 := \frac{J_5}{J_2 \cdot J_3}, \quad t_4 := \frac{J_6}{J_2 \cdot J_4}, \quad t_5 := \frac{J_7}{J_2 \cdot J_5}, \quad t_6 := \frac{J_8}{J_2^4},$$

There is an algebraic relation

$$(12) \quad T(t_1, \dots, t_6) = 0$$

that such invariants satisfy, computed in [27]. The field of invariants \mathcal{S}_8 of binary octavics is $\mathcal{S}_8 = k(t_1, \dots, t_6)$, where t_1, \dots, t_6 satisfies the equation $T(t_1, \dots, t_6) = 0$. Hence, we have an explicit description of the hyperelliptic moduli \mathcal{H}_3 ; see [27] for details.

Throughout this paper we will use the following important result

Lemma 5 (Shaska [27]). *Two genus 3 hyperelliptic curves C and C' , defined over an algebraically closed field k of characteristic zero, with J_2, J_3, J_4, J_5 nonzero are isomorphic over k if and only if*

$$t_i(C) = t_i(C'), \quad \text{for } i = 1, \dots, 6.$$

In the cases of curves when t_1, \dots, t_6 are not defined we will define new invariants as suggested in [27]. From [27, Lemma 4] we know that J_2, \dots, J_7 can't all be 0, otherwise the binary form would have a multiple root.

To describe the moduli points in cases when absolute invariants are not defined is not difficult. In this case, one has to treat each case separately when any of the invariants J_2, \dots, J_5 are zero. Indeed, we can define invariants depending of which of the invariants is nonzero.

If $J_2 \neq 0$, then we define

$$(13) \quad i_1 = \frac{J_3^2}{J_2^3}, \quad i_2 = \frac{J_4}{J_2^2}, \quad i_3 = \frac{J_5^2}{J_2^5}, \quad i_4 = \frac{J_6}{J_2^3}, \quad i_5 = \frac{J_7^2}{J_2^7}, \quad i_6 = \frac{J_8}{J_2^4}$$

If $J_2 = 0$ then we pick the smallest degree invariant among J_3, \dots, J_7 which is not zero. Let $J_2 = 0$ and $J_3 \neq 0$. Define

$$h_1 = \frac{J_4^3}{J_3^4}, \quad h_2 = \frac{J_5^3}{J_3^5}, \quad h_3 = \frac{J_6}{J_3^2}, \quad h_4 = \frac{J_7^3}{J_3^7}, \quad h_5 = \frac{J_8^3}{J_3^8}$$

Let $J_2 = J_3 = 0$ and $J_4 \neq 0$. Then we have

$$j_1 = \frac{J_5^4}{J_4^5}, \quad j_2 = \frac{J_6^2}{J_4^3}, \quad j_3 = \frac{J_7^4}{J_4^7}, \quad j_4 = \frac{J_8}{J_4^2}$$

Let $J_2 = J_3 = J_4 = 0$ and $J_5 \neq 0$. Then

$$k_1 = \frac{J_6^5}{J_5^6}, \quad k_2 = \frac{J_7^5}{J_5^7}, \quad k_3 = \frac{J_8^5}{J_5^8}$$

Let us assume that $J_2 = J_3 = J_4 = J_5 = 0$. In this case, we define the absolute invariants

$$\tau_1 := \frac{J_7^6}{J_6^7}, \quad \tau_2 = \frac{J_8^3}{J_6^4}$$

There is only one curve in the case when $J_2 = \dots = J_6 = 0$, namely $Y^2 = X^7 - 1$. In our discussion in section 5 we will see cases when $J_3 = J_5 = J_7 = 0$. In this case we will use invariants defined in Eq. (13).

Since a tuple (t_1, \dots, t_6) uniquely determines the isomorphism class of a curve then we will study the locus of the curves with a fixed automorphism group G in terms of such invariants (t_1, \dots, t_6) . The only interesting cases are groups G which have non-hyperelliptic involutions; see [15] or [4].

To make it easier to state some of the results in the following sections we define the absolute invariants of \mathcal{X} as follows

$$\mathbf{p}(\mathcal{X}) = \begin{cases} (t_1, \dots, t_6) & \text{if } J_2, \dots, J_5 \text{ are nonzero} \\ (i_1, \dots, i_6) & \text{if } J_2 \neq 0 \wedge (J_3 = 0 \vee J_4 = 0 \vee J_5 = 0) \\ (h_1, \dots, h_5) & \text{if } J_2 = 0 \wedge J_3 \neq 0 \\ (j_1, j_2, j_3, j_4) & \text{if } J_2 = J_3 = 0 \wedge J_4 \neq 0 \\ (k_1, k_2, k_3) & \text{if } J_2 = J_3 = J_4 = 0 \wedge J_5 \neq 0 \\ (\tau_1, \tau_2) & \text{if } J_2 = J_3 = J_4 = J_5 = 0 \end{cases}$$

For each case of the above we determine the equation of the moduli space \mathcal{H}_3 . For the first case this equation is Eq. (12). The rest of the cases are described briefly below. The $\mathbf{p}(\mathcal{X})$ determines uniquely the isomorphism class of \mathcal{X} .

Computational benefits of these invariants are that they are of small degree and therefore nicer especially when we deal with families of curves and have to compute symbolically. In each case for $\mathbf{p}(\mathcal{X})$ we can compute the equation of the moduli in terms of the corresponding invariants analogous to the $T(t_1, \dots, t_6) = 0$ as in Eq. (12).

4.1. Computing the locus \mathcal{S} . Let \mathcal{S} denote the locus of genus 3 hyperelliptic curves with elliptic involutions. It follows from the theory of Hurwitz spaces that this is an irreducible 3-dimensional subvariety of \mathcal{H}_3 ; see [15, 24]. In [15] it is shown that $k(\mathcal{S}) = k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$. In this paper, we will give an explicit computational proof of this result and provide a birational parametrization of the locus \mathcal{S} . We will outline all the computations and display only those results which are reasonable to display in this paper.

Every genus 3 hyperelliptic curve which has an elliptic involution is isomorphic to a curve with equation as in Eq. (4). The obvious strategy would be to compute the invariants t_1, \dots, t_6 in terms of a, b, c and then eliminate a, b, c from these equations. This is rather difficult computationally. Since dihedral invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ are invariant under coordinate changes in $\mathbb{P}^1(k)^1$ then we can express t_1, \dots, t_6 in terms of such invariants as stated in Theorem 1. In the next few paragraphs we describe these computations.

Let \mathcal{X}_3 be a genus 3 hyperelliptic curve with equation as in Eq. (4). Notice that from the definitions of the dihedral invariants in Eq. (5) we have

$$(14) \quad b = \frac{\mathfrak{s}_3}{a^2 + c^2}, \quad b^2 = \frac{\mathfrak{s}_3^2}{\mathfrak{s}_4 + 2\mathfrak{s}_2^2}, \quad a^8 + c^8 = \mathfrak{s}_4^2 - 2\mathfrak{s}_2^4, \quad (a^2 + c^2)^2 = \mathfrak{s}_4 + 2\mathfrak{s}_2^2$$

We denote by $\lambda := a^2 + c^2$. Then, $\lambda^2 = \mathfrak{s}_4 + 2\mathfrak{s}_2^2$ and $\lambda a^2 = (a^2 + c^2)a^2 = a^4 + \mathfrak{s}_2^2$. By changing the coordinate by

$$X \rightarrow (a^2 + c^2) \sqrt{a} X$$

we get the curve

$$Y^2 = \lambda^8 a^4 X^8 + \lambda^6 a^4 X^6 + \lambda^4 b a^2 X^4 + \lambda^2 c a X^2 + 1$$

Notice that the coefficient of X^4 is

$$\lambda^4 b a^2 = (a^2 + c^2)^2 \cdot \lambda^2 b a^2 = b(a^2 + c^2) \cdot a^2(a^2 + c^2) \cdot \lambda^2 = \mathfrak{s}_3 \cdot (a^4 + \mathfrak{s}_2) \cdot \lambda^2$$

Then, we have the curve with equation

$$(15) \quad Y^2 = A X^8 + \frac{A}{\mathfrak{s}_4 + 2\mathfrak{s}_2^2} X^6 + \frac{\mathfrak{s}_3(A + \mathfrak{s}_2^2)}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^3} X^4 + \frac{\mathfrak{s}_2}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^3} X^2 + \frac{1}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^4}$$

where $A = a^4$. Notice that by substituting $c = \frac{\mathfrak{s}_2}{a}$ in the definition of \mathfrak{s}_4 we get that $A + \frac{\mathfrak{s}_2^4}{A} = \mathfrak{s}_4$, which says that A satisfies the equation

$$(16) \quad A^2 - \mathfrak{s}_4 A + \mathfrak{s}_2^4 = 0$$

We will see how this equation will be useful when discussing the field of definition of the curve \mathcal{X}_3 .

From the Eq. (4) we compute the invariants J_2, \dots, J_8 . By performing the above substitutions we get the following expressions

(17)

$$J_2 = 2(140 \mathfrak{s}_4 + 280 \mathfrak{s}_2^2 + 5 \mathfrak{s}_2 \mathfrak{s}_4 + 10 \mathfrak{s}_2^3 + \mathfrak{s}_3^2)$$

$$J_3 = 2(6 \mathfrak{s}_3^3 + 525 \mathfrak{s}_4^2 + 2100 \mathfrak{s}_4 \mathfrak{s}_2^2 + 2100 \mathfrak{s}_2^4 - 55 \mathfrak{s}_2 \mathfrak{s}_3 \mathfrak{s}_4 - 110 \mathfrak{s}_2^3 \mathfrak{s}_3 + 1960 \mathfrak{s}_3 \mathfrak{s}_4 + 3920 \mathfrak{s}_3 \mathfrak{s}_2^2)$$

$$J_4 = 2^6 (\mathfrak{s}_3^4 + 126 \mathfrak{s}_3 \mathfrak{s}_4^2 + 504 \mathfrak{s}_3 \mathfrak{s}_4 \mathfrak{s}_2^2 + 504 \mathfrak{s}_3 \mathfrak{s}_2^4 + 38416 \mathfrak{s}_4^2 + 153664 \mathfrak{s}_4 \mathfrak{s}_2^2 + 153664 \mathfrak{s}_2^4 \\ - 784 \mathfrak{s}_2^3 \mathfrak{s}_4 - 784 \mathfrak{s}_2^5 + 4 \mathfrak{s}_4^2 \mathfrak{s}_2^2 + 16 \mathfrak{s}_4 \mathfrak{s}_2^4 + 16 \mathfrak{s}_2^6 - 392 \mathfrak{s}_4 \mathfrak{s}_3^2 - 784 \mathfrak{s}_2^2 \mathfrak{s}_3^2 + 31 \mathfrak{s}_2 \mathfrak{s}_4 \mathfrak{s}_3^2 \\ - 196 \mathfrak{s}_2 \mathfrak{s}_4^2 + 62 \mathfrak{s}_2^3 \mathfrak{s}_3^2)$$

$$J_5 = 2^5 (76832 \mathfrak{s}_3 \mathfrak{s}_4^2 + 307328 \mathfrak{s}_3 \mathfrak{s}_2^4 + 123480 \mathfrak{s}_4^2 \mathfrak{s}_2^2 + 246960 \mathfrak{s}_4 \mathfrak{s}_2^4 + 1148 \mathfrak{s}_2^4 \mathfrak{s}_3^2 + 287 \mathfrak{s}_4^2 \mathfrak{s}_3^2 \\ - 1568 \mathfrak{s}_3^3 \mathfrak{s}_2^2 + 41552 \mathfrak{s}_2^5 \mathfrak{s}_3 - 26 \mathfrak{s}_2^3 \mathfrak{s}_3^3 - 1680 \mathfrak{s}_2^5 \mathfrak{s}_4 - 208 \mathfrak{s}_3 \mathfrak{s}_2^6 - 140 \mathfrak{s}_2 \mathfrak{s}_4^3 - 840 \mathfrak{s}_2^3 \mathfrak{s}_4^2 \\ + 20580 \mathfrak{s}_4^3 + 2 \mathfrak{s}_3^5 - 1120 \mathfrak{s}_2^7 + 307328 \mathfrak{s}_3 \mathfrak{s}_4 \mathfrak{s}_2^2 + 10388 \mathfrak{s}_2 \mathfrak{s}_3 \mathfrak{s}_4^2 + 41552 \mathfrak{s}_2^3 \mathfrak{s}_3 \mathfrak{s}_4 \\ - 52 \mathfrak{s}_3 \mathfrak{s}_4^2 \mathfrak{s}_2^2 + 164640 \mathfrak{s}_2^6 - 784 \mathfrak{s}_3^3 \mathfrak{s}_4 - 208 \mathfrak{s}_3 \mathfrak{s}_4 \mathfrak{s}_2^4 + 1148 \mathfrak{s}_2^2 \mathfrak{s}_3^2 \mathfrak{s}_4 - 13 \mathfrak{s}_2 \mathfrak{s}_3^3 \mathfrak{s}_4)$$

$$J_6 = 2^9 (2 \mathfrak{s}_2 \mathfrak{s}_4 + 4 \mathfrak{s}_2^3 - 196 \mathfrak{s}_4 - 392 \mathfrak{s}_2^2 + \mathfrak{s}_3^2) (\mathfrak{s}_3^4 - 378 \mathfrak{s}_3 \mathfrak{s}_4^2 - 1512 \mathfrak{s}_3 \mathfrak{s}_4 \mathfrak{s}_2^2 - 1512 \mathfrak{s}_3 \mathfrak{s}_2^4 \\ - 10192 \mathfrak{s}_2^3 \mathfrak{s}_4 - 10192 \mathfrak{s}_2^5 - 77 \mathfrak{s}_2 \mathfrak{s}_4 \mathfrak{s}_3^2 - 154 \mathfrak{s}_2^3 \mathfrak{s}_3^2 + 4 \mathfrak{s}_4^2 \mathfrak{s}_2^2 + 16 \mathfrak{s}_4 \mathfrak{s}_2^4 + 16 \mathfrak{s}_2^6 \\ + 38416 \mathfrak{s}_4^2 + 153664 \mathfrak{s}_4 \mathfrak{s}_2^2 + 153664 \mathfrak{s}_2^4 - 392 \mathfrak{s}_4 \mathfrak{s}_3^2 - 784 \mathfrak{s}_2^2 \mathfrak{s}_3^2 - 2548 \mathfrak{s}_2 \mathfrak{s}_4^2)$$

$$J_7 = 2^7 (1120 \mathfrak{s}_2^4 \mathfrak{s}_4^3 - 120472576 \mathfrak{s}_3 \mathfrak{s}_2^6 - 34300 \mathfrak{s}_2 \mathfrak{s}_4^4 + 3360 \mathfrak{s}_2^6 \mathfrak{s}_4^2 + 4480 \mathfrak{s}_2^8 \mathfrak{s}_4 + 140 \mathfrak{s}_4^4 \mathfrak{s}_2^2 \\ - 203840 \mathfrak{s}_3 \mathfrak{s}_2^8 + 608 \mathfrak{s}_3 \mathfrak{s}_2^9 + 4410 \mathfrak{s}_3 \mathfrak{s}_4^4 - 39200 \mathfrak{s}_2^5 \mathfrak{s}_3^3 - 931 \mathfrak{s}_3^4 \mathfrak{s}_4^2 - 3724 \mathfrak{s}_3^4 \mathfrak{s}_2^4 \\ - 90 \mathfrak{s}_3^5 \mathfrak{s}_2^3 - 1176 \mathfrak{s}_3^5 \mathfrak{s}_4 - 2352 \mathfrak{s}_3^5 \mathfrak{s}_2^2 + 161896 \mathfrak{s}_3^2 \mathfrak{s}_4^3 + 8344 \mathfrak{s}_2^7 \mathfrak{s}_3^2 - 274400 \mathfrak{s}_2^3 \mathfrak{s}_4^3 \\ + 230496 \mathfrak{s}_3^3 \mathfrak{s}_4^2 + 921984 \mathfrak{s}_3^3 \mathfrak{s}_2^4 + 1295168 \mathfrak{s}_2^6 \mathfrak{s}_3^2 + 129077760 \mathfrak{s}_2^6 \mathfrak{s}_4 - 1097600 \mathfrak{s}_2^7 \mathfrak{s}_4 \\ - 15059072 \mathfrak{s}_3 \mathfrak{s}_4^3 + 96808320 \mathfrak{s}_2^4 \mathfrak{s}_4^2 + 29196160 \mathfrak{s}_2^7 \mathfrak{s}_3 + 2240 \mathfrak{s}_2^{10} + 4033680 \mathfrak{s}_4^4 \\ + 2 \mathfrak{s}_3^7 - 90354432 \mathfrak{s}_3 \mathfrak{s}_4^2 \mathfrak{s}_2^2 - 180708864 \mathfrak{s}_3 \mathfrak{s}_4 \mathfrak{s}_2^4 - 270480 \mathfrak{s}_3 \mathfrak{s}_2^6 \mathfrak{s}_4 - 45 \mathfrak{s}_3^5 \mathfrak{s}_2 \mathfrak{s}_4 \\ + 6258 \mathfrak{s}_2^3 \mathfrak{s}_3^2 \mathfrak{s}_4^2 + 912 \mathfrak{s}_3 \mathfrak{s}_2^7 \mathfrak{s}_4 + 971376 \mathfrak{s}_2^2 \mathfrak{s}_3^2 \mathfrak{s}_4^2 - 9800 \mathfrak{s}_2 \mathfrak{s}_3^3 \mathfrak{s}_4^2 + 345 \mathfrak{s}_2^2 \mathfrak{s}_3^3 \mathfrak{s}_4^2 \\ + 456 \mathfrak{s}_3 \mathfrak{s}_2^5 \mathfrak{s}_4^2 - 39200 \mathfrak{s}_2^3 \mathfrak{s}_3^3 \mathfrak{s}_4 + 64538880 \mathfrak{s}_2^8 + 1380 \mathfrak{s}_2^6 \mathfrak{s}_3^3 \\ - 3724 \mathfrak{s}_3^4 \mathfrak{s}_4 \mathfrak{s}_2^2 + 921984 \mathfrak{s}_3^3 \mathfrak{s}_4 \mathfrak{s}_2^2 + 1043 \mathfrak{s}_2 \mathfrak{s}_3^2 \mathfrak{s}_4^3 + 21897120 \mathfrak{s}_2^3 \mathfrak{s}_3 \mathfrak{s}_4^2 + 76 \mathfrak{s}_3 \mathfrak{s}_2^3 \mathfrak{s}_4^3 \\ + 43794240 \mathfrak{s}_2^5 \mathfrak{s}_3 \mathfrak{s}_4 + 1380 \mathfrak{s}_2^4 \mathfrak{s}_3^3 \mathfrak{s}_4 + 3649520 \mathfrak{s}_2 \mathfrak{s}_3 \mathfrak{s}_4^3 + 1942752 \mathfrak{s}_2^4 \mathfrak{s}_3^2 \mathfrak{s}_4 + 980 \mathfrak{s}_3 \mathfrak{s}_4^3 \mathfrak{s}_2^2 \\ + 32269440 \mathfrak{s}_4^3 \mathfrak{s}_2^2 - 99960 \mathfrak{s}_3 \mathfrak{s}_2^4 \mathfrak{s}_4^2 - 548800 \mathfrak{s}_2^9 + 12516 \mathfrak{s}_2^5 \mathfrak{s}_3^2 \mathfrak{s}_4 - 823200 \mathfrak{s}_2^5 \mathfrak{s}_4^2)$$

We do not display J_8 but it is easy to compute as the previous ones. The reader who want to obtain the above expressions can use computational packages. For example, "algsb" would work in Maple and similar commands in other packages.

Hence, we can write now t_1, \dots, t_6 in terms of the dihedral invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$

$$(18)$$

$$t_1 = \frac{1}{2} \cdot \frac{(3920\mathfrak{s}_3\mathfrak{s}_2^2 + 2100\mathfrak{s}_4^2 + 2100\mathfrak{s}_2^2\mathfrak{s}_4 + 525\mathfrak{s}_2^4 + 7840\mathfrak{s}_3\mathfrak{s}_4 + 24\mathfrak{s}_3^3 - 110\mathfrak{s}_3\mathfrak{s}_4\mathfrak{s}_2 - 55\mathfrak{s}_3\mathfrak{s}_2^3)^2(2\mathfrak{s}_4 + \mathfrak{s}_2^2)^2}{(560\mathfrak{s}_4 + 280\mathfrak{s}_2^2 + 10\mathfrak{s}_2\mathfrak{s}_4 + 5\mathfrak{s}_2^3 + 4\mathfrak{s}_3^2)^3}$$

$$t_2 = \frac{64}{(560\mathfrak{s}_4 + 280\mathfrak{s}_2^2 + 10\mathfrak{s}_2\mathfrak{s}_4 + 5\mathfrak{s}_2^3 + 4\mathfrak{s}_3^2)^2} (38416\mathfrak{s}_2^4 + \mathfrak{s}_2^6 + 4\mathfrak{s}_4^2\mathfrak{s}_2^2 + 4\mathfrak{s}_2^4\mathfrak{s}_4 - 392\mathfrak{s}_2^3\mathfrak{s}_4 + 153664\mathfrak{s}_2^2\mathfrak{s}_4 - 392\mathfrak{s}_4^2\mathfrak{s}_2$$

$$+ 4\mathfrak{s}_3^4 - 98\mathfrak{s}_2^5 + 504\mathfrak{s}_3\mathfrak{s}_2^2\mathfrak{s}_4 - 1568\mathfrak{s}_3^2\mathfrak{s}_4 - 784\mathfrak{s}_3^2\mathfrak{s}_2^2 + 504\mathfrak{s}_3\mathfrak{s}_4^2 + 126\mathfrak{s}_3\mathfrak{s}_2^4 + 62\mathfrak{s}_3^2\mathfrak{s}_4\mathfrak{s}_2 + 153664\mathfrak{s}_4^2 + 31\mathfrak{s}_3^2\mathfrak{s}_2^3)$$

$$t_3 = -\frac{32}{(560\mathfrak{s}_4 + 280\mathfrak{s}_2^2 + 10\mathfrak{s}_2\mathfrak{s}_4 + 5\mathfrak{s}_2^3 + 4\mathfrak{s}_3^2)} \cdot \frac{M}{N}$$

$$M = -123480\mathfrak{s}_4^2\mathfrak{s}_2^2 - 61740\mathfrak{s}_2^4\mathfrak{s}_4 - 307328\mathfrak{s}_3\mathfrak{s}_4^2 - 76832\mathfrak{s}_3\mathfrak{s}_2^4 - 5194\mathfrak{s}_3\mathfrak{s}_2^5 + 13\mathfrak{s}_3^3\mathfrak{s}_2^3 + 3136\mathfrak{s}_3^3\mathfrak{s}_4 + 1568\mathfrak{s}_3^3\mathfrak{s}_2^2$$

$$- 1148\mathfrak{s}_3^2\mathfrak{s}_4^2 + 280\mathfrak{s}_4^3\mathfrak{s}_2 + 420\mathfrak{s}_2^3\mathfrak{s}_4^2 + 210\mathfrak{s}_2^5\mathfrak{s}_4 + 13\mathfrak{s}_3\mathfrak{s}_2^6 - 287\mathfrak{s}_3^2\mathfrak{s}_2^4 - 10290\mathfrak{s}_2^6 - 8\mathfrak{s}_3^5 - 82320\mathfrak{s}_4^3 + 35\mathfrak{s}_2^7$$

$$- 307328\mathfrak{s}_3\mathfrak{s}_2^2\mathfrak{s}_4 - 20776\mathfrak{s}_3\mathfrak{s}_4^2\mathfrak{s}_2 + 52\mathfrak{s}_3\mathfrak{s}_4^2\mathfrak{s}_2^2 - 20776\mathfrak{s}_3\mathfrak{s}_2^3\mathfrak{s}_4 + 52\mathfrak{s}_3\mathfrak{s}_2^4\mathfrak{s}_4 + 26\mathfrak{s}_3^3\mathfrak{s}_4\mathfrak{s}_2 - 1148\mathfrak{s}_3^2\mathfrak{s}_4\mathfrak{s}_2^2$$

$$N = 3920\mathfrak{s}_3\mathfrak{s}_2^2 + 2100\mathfrak{s}_4^2 + 2100\mathfrak{s}_2^2\mathfrak{s}_4 + 525\mathfrak{s}_2^4 + 7840\mathfrak{s}_3\mathfrak{s}_4 + 24\mathfrak{s}_3^3 - 110\mathfrak{s}_3\mathfrak{s}_4\mathfrak{s}_2 - 55\mathfrak{s}_3\mathfrak{s}_2^3$$

$$t_4 = -\frac{A}{B} \cdot \frac{(8(\mathfrak{s}_2^3 + 2\mathfrak{s}_2\mathfrak{s}_4 - 392\mathfrak{s}_4 - 196\mathfrak{s}_2^2 + 2\mathfrak{s}_3^2))}{(560\mathfrak{s}_4 + 280\mathfrak{s}_2^2 + 10\mathfrak{s}_2\mathfrak{s}_4 + 5\mathfrak{s}_2^3 + 4\mathfrak{s}_3^2)}$$

$$A = -38416\mathfrak{s}_2^4 - \mathfrak{s}_2^6 - 4\mathfrak{s}_4^2\mathfrak{s}_2^2 - 4\mathfrak{s}_2^4\mathfrak{s}_4 + 5096\mathfrak{s}_2^3\mathfrak{s}_4 - 153664\mathfrak{s}_2^2\mathfrak{s}_4 + 5096\mathfrak{s}_4^2\mathfrak{s}_2 - 4\mathfrak{s}_3^4 + 1274\mathfrak{s}_2^5 + 1512\mathfrak{s}_3\mathfrak{s}_2^2\mathfrak{s}_4$$

$$+ 1568\mathfrak{s}_3^2\mathfrak{s}_4 + 784\mathfrak{s}_3^2\mathfrak{s}_2^2 + 1512\mathfrak{s}_3\mathfrak{s}_4^2 + 378\mathfrak{s}_3\mathfrak{s}_2^4 + 154\mathfrak{s}_3^2\mathfrak{s}_4\mathfrak{s}_2 - 153664\mathfrak{s}_4^2 + 77\mathfrak{s}_3^2\mathfrak{s}_2^3)$$

$$B = 38416\mathfrak{s}_2^4 + \mathfrak{s}_2^6 + 4\mathfrak{s}_4^2\mathfrak{s}_2^2 + 4\mathfrak{s}_2^4\mathfrak{s}_4 - 392\mathfrak{s}_2^3\mathfrak{s}_4 + 153664\mathfrak{s}_2^2\mathfrak{s}_4 - 392\mathfrak{s}_4^2\mathfrak{s}_2 + 4\mathfrak{s}_3^4 - 98\mathfrak{s}_2^5 + 504\mathfrak{s}_3\mathfrak{s}_2^2\mathfrak{s}_4$$

$$- 1568\mathfrak{s}_3^2\mathfrak{s}_4 - 784\mathfrak{s}_3^2\mathfrak{s}_2^2 + 504\mathfrak{s}_3\mathfrak{s}_4^2 + 126\mathfrak{s}_3\mathfrak{s}_2^4 + 62\mathfrak{s}_3^2\mathfrak{s}_4\mathfrak{s}_2 + 153664\mathfrak{s}_4^2 + 31\mathfrak{s}_3^2\mathfrak{s}_2^3$$

We are not displaying t_5 and t_6 .

Hence, we have a map give by the above equations

$$k^3 \setminus \{\Delta = 0\} \rightarrow \mathcal{S} = \mathcal{M}_3^b \cap \mathcal{H}_3$$

$$(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) \rightarrow (t_1, t_2, t_3, t_4, t_5, t_6)$$

which as it will be shown in the next theorem is birational.

Theorem 2. $k(\mathcal{S}) = k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$.

Proof. Since $k(\mathcal{S})$ is a subfield of $k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$ which contains all $k(t_i)$ for $i = 1, \dots, 6$ then $[k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) : k(\mathcal{S})]$ must divide each of degrees if t_i . The degrees of t_i as rational functions in $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ are respectively 12, 6, 7, 9, 10, 12. Hence, $[k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) : k(\mathcal{S})] = 1$. This completes the proof. \square

This was also proved in [15] for any genus $g \geq 2$. Here we provide a direct computational proof and explicitly determine the formulas for $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ as rational functions in terms of t_1, \dots, t_6 . We have the following theorem.

Theorem 3. *The space $\mathcal{S} := \mathcal{M}_3^b \cap \mathcal{H}_3$ is an irreducible, codimension 1, rational subvariety of \mathcal{M}_3^b . Its defining equations are*

$$(19) \quad F_i(J_2, \dots, J_8) = 0, \quad i = 1 \dots 5$$

as displayed in [29]. The map

$$(20) \quad k^3 \setminus \{\Delta = 0\} \rightarrow \mathcal{S} := \mathcal{M}_3^b \cap \mathcal{H}_3$$

$$(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) \rightarrow (t_1, \dots, t_6)$$

given by Eq. (18) (in homogeneous coordinates by the formulas (17)) is birational and surjective.

Proof. From Theorem 1, ii) we have that $\mathcal{M}_3^b \cap \mathcal{H}_3$ is birationally isomorphic to the coarse moduli space of smooth curves of genus 2 together with a nontrivial divisor class of order 2. Since this space is an irreducible, 3-dimensional, rational variety then the first part of the theorem is proved.

It remains to show the map in the Theorem is birational. We need to show that the degree of the field extension $k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)/k(\mathcal{S})$ is 1. For this we use the functions $t_1, t_2, t_3, t_4, t_5, t_6$ in $k(\mathcal{H}_3)$. The fact that $[k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) : k(\mathcal{S})] = 1$ comes straight from the computation of the locus \mathcal{S} where we get rational expressions for the $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ or from Theorem 2. This completes the proof. \square

4.2. Field of moduli vs field of definition. It is a classical problem in the arithmetic of the algebraic curves to try to find an equation of the curve in terms of the moduli point corresponding to this curve. In other words, this means that given the moduli point $\mathfrak{p}(\mathcal{X})$, could we determine an equation for \mathcal{X} in terms of the coordinates of $\mathfrak{p}(\mathcal{X})$. If an equation of the curve can be found in terms of coordinates of the moduli point we say that the field of moduli is the same with the minimal field of definition. In 2003 the first author conjectured that this would be the case for all hyperelliptic curves with extra involutions [23, 25]. It is true for $g = 2$ and as we will see next it is true for all genus 3 hyperelliptic curves in $\mathfrak{p} \in \mathcal{M}_3^b \cap \mathcal{H}_3$ such that $|\text{Aut}(\mathfrak{p})| > 2$. There have been claims on whether the above conjecture is true or false and some confusion from work of Huggins [17] and Fuertes [12] which seem to come from different definitions of the field of moduli.

Summarizing we have the results of section 4.1 and Eq. (15) and (16) we have the following:

Proposition 2. *Let $[\mathcal{X}] \in \mathcal{M}_3^b \cap \mathcal{H}_3$. Then the following hold true:*

i) \mathcal{X} is isomorphic to a curve with equation

$$Y^2 = AX^8 + \frac{A}{\mathfrak{s}_4 + 2\mathfrak{s}_2^2} X^6 + \frac{\mathfrak{s}_3(A + \mathfrak{s}_2^2)}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^3} X^4 + \frac{\mathfrak{s}_2}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^3} X^2 + \frac{1}{(\mathfrak{s}_4 + 2\mathfrak{s}_2^2)^4}$$

where A satisfies

$$(21) \quad A^2 - \mathfrak{s}_4 A + \mathfrak{s}_2^4 = 0,$$

for some $(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4 \in k^3 \setminus \{\Delta_{\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4} = 0\})$ and $\Delta_{\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4}$ as in Eq. (6).

ii) Let F denote the field of moduli of \mathcal{X} and F' its minimal field of definition. Then, $F' \subset F(A)$ and $[F' : F] \leq 2$. An equation of \mathcal{X} over $F(A)$ is given by Eq. (21).

iii) If the discriminant $d = \mathfrak{s}_4^2 - 4\mathfrak{s}_2^4$ of the quadratic in Eq. (21) is a complete square in $k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$ then the corresponding curve is defined over its field of moduli.

Proof. Part i) was proved in section 4.1. The field of moduli of a curve \mathcal{X} is $F = k(t_1, \dots, t_6)$. Hence, from Theorem 2, $F = k(\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4)$. Since \mathcal{X} is defined via Eq. (21) over $F(\sqrt{\mathfrak{s}_4^2 - 4\mathfrak{s}_2^4})$, then $F' \subset F(A)$ and $[F' : F] \leq 2$. Part iii) is clearly true. \square

The above result improves on the bound of the degree of $[F' : F] \leq 8$ as shown in [18]. It is expected that the field of moduli is a field of definition for all curves $\mathcal{X} \in \mathcal{M}_3^b \cap \mathcal{H}_3$. A generalization of methods for $g = 2$ should provide an algorithm also for $g = 3$.

5. SINGULAR LOCUS OF \mathcal{S} , CLASSIFICATION OF STRATA OF THE HYPERELLIPTIC MODULI

In this section we give a classification of the strata of the hyperelliptic moduli 3. The stratum of the hyperelliptic moduli has been known to the classical algebraic geometers. Indeed, it is the only case that was considered fully known even though explicit descriptions were not available even for small genus (i.e., $g = 2$). On the turn of the new century a couple of papers appeared for the case of genus $g = 2$; see [13] and [30].

5.1. Singular loci in terms of dihedral invariants. Next, we will characterize each one of these loci in terms of the \mathfrak{s} -invariants. The proof of the following theorem can be found in [15], where such relations were determined by studying the group action on the invariants $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$. Here we give a more computational proof.

5.2. 2-dimensional strata. There are two 2-dimensional loci in \mathcal{H}_3 which correspond to the case when the reduced automorphism group of the curve is isomorphic to V_4 . Indeed, the following is true for any genus $g \geq 2$; see [16, Theorem 3].

Let \mathcal{X}_g be a genus g hyperelliptic curve with an extra involution, $\overline{\text{Aut}}(\mathcal{X}_g)$ its reduced automorphism group, and $(\mathfrak{s}_4, \dots, \mathfrak{s}_g)$ its corresponding dihedral invariants as defined in [16].

If $V_4 \hookrightarrow \overline{\text{Aut}}(\mathcal{X}_g)$ then $2^{g-1} \mathfrak{s}_1^2 = \mathfrak{s}_g^{g+1}$. Moreover, if g is odd then $V_4 \hookrightarrow \overline{\text{Aut}}(\mathcal{X}_g)$ implies that

$$(22) \quad (2^r \mathfrak{s}_1 - \mathfrak{s}_g^{r+1}) (2^r \mathfrak{s}_1 + \mathfrak{s}_g^{r+1}) = 0$$

where $r = \lfloor \frac{g-1}{2} \rfloor$. The first factor corresponds to the case when involutions of $V_4 \hookrightarrow \overline{\text{Aut}}(\mathcal{X}_g)$ lift to involutions in $\text{Aut}(\mathcal{X}_g)$, the second factor corresponds to the case when two of the involutions of $V_4 \hookrightarrow \overline{\text{Aut}}(\mathcal{X}_g)$ lift to elements of order 4 in $\text{Aut}(\mathcal{X}_g)$.

In the case of genus $g = 3$, we have $\mathfrak{s}_1 = \mathfrak{s}_4$ and $\mathfrak{s}_g = \frac{\mathfrak{s}_2}{2}$ and the relation becomes $\mathfrak{s}_4 - 2\mathfrak{s}_2^2 = 0$. This gives exactly the cases when the full automorphism group is \mathbb{Z}_2^3 . We will verify such fact directly below.

5.2.1. The automorphism group is isomorphic to $G \cong \mathbb{Z}_2^3$: The equation of the curve from Table 3 of [20] is

$$y^2 = (x^4 + ax^2 + 1)(x^4 + bx^2 + 1)$$

Let new parameters u and v be as follows

$$u := a + b, \quad \text{and} \quad v = ab.$$

Then we have

$$(23) \quad y^2 = x^8 + ux^6 + (v+2)x^4 + ux^2 + 1$$

The dihedral invariants are

$$\mathfrak{s}_4 = 2u^4, \quad \mathfrak{s}_3 = 2u^2(v+2), \quad \mathfrak{s}_2 = u^2$$

Then directly we can verify $u^2 = \mathfrak{s}_2$, $v = \frac{\mathfrak{s}_3 - 2\mathfrak{s}_2}{2\mathfrak{s}_2}$ and

$$(24) \quad \mathfrak{s}_4 - 2\mathfrak{s}_2^2 = 0.$$

By transforming the coordinate X as $X \rightarrow \sqrt{u}X$ on the curve in Eq.(23) we get

$$Y^2 = u^4 X^8 + u^4 X^6 + (v+2)(u^2)X^4 + u^2 X^2 + 1$$

or

$$(25) \quad Y^2 = \mathfrak{s}_2^2 X^8 + \mathfrak{s}_2^2 X^6 + \frac{\mathfrak{s}_3}{2} X^4 + \mathfrak{s}_2 X^2 + 1$$

We compute the invariants t_1, \dots, t_6 in terms of $\mathfrak{s}_3, \mathfrak{s}_2$. Eliminating \mathfrak{s}_3 and \mathfrak{s}_2 from the system of equations gives the locus $\mathcal{S}(\mathbb{Z}_2^3)$ and rational expressions of $\mathfrak{s}_3, \mathfrak{s}_2$ in terms of t_1, \dots, t_6 . The computations are long and the results involve very large expressions. Instead, we provide a quicker proof for the reader which is easier to check.

From the expressions of t_1, \dots, t_6 in terms of $\mathfrak{s}_3, \mathfrak{s}_2$ we eliminate \mathfrak{s}_3 . In other words, \mathfrak{s}_3 is easily written as a rational function in terms of $\mathfrak{s}_2, t_1, \dots, t_6$. Any software package such as maple or Mathematica will be able to do this. We are left with 5 equations of degree 20, 18, 16, 23, and 21. Since the function field $k(\mathcal{S}(\mathbb{Z}_2^3))$ is a subfield of $k(\mathfrak{s}_2)$ then the degree of this extension $[k(\mathfrak{s}_2), k(\mathcal{S}(\mathbb{Z}_2^3))]$ is a common divisor of 20, 18, 16, 23, and 21. Therefore, $[k(\mathfrak{s}_2), k(\mathcal{S}(\mathbb{Z}_2^3))] = 1$ and $k(\mathcal{S}(\mathbb{Z}_2^3)) = k(\mathfrak{s}_2)$.

Lemma 6. *Every genus 3 hyperelliptic curve with full automorphism group isomorphic to \mathbb{Z}_2^3 has equation*

$$(26) \quad Y^2 = \mathfrak{s}_2^2 X^8 + \mathfrak{s}_2^2 X^6 + \frac{\mathfrak{s}_3}{2} X^4 + \mathfrak{s}_2 X^2 + 1$$

for $\mathfrak{s}_3, \mathfrak{s}_2 \neq 0, 4$. Moreover, $k(\mathcal{S}(\mathbb{Z}_2^3)) = k(\mathfrak{s}_2, \mathfrak{s}_3)$ and therefore the field of moduli is a field of definition.

The result of the Lemma above can be obtained directly by the V_4 -locus, by enforcing the equation $2\mathfrak{s}_4 - \mathfrak{s}_2^2 = 0$. The expressions for $\mathfrak{s}_3, \mathfrak{s}_2$ and the equations of $\mathcal{S}(\mathbb{Z}_2^3)$ in terms of t_1, \dots, t_6 are displayed in [29].

5.2.2. *The automorphism group is isomorphic to $G \cong \mathbb{Z}_4$:* This is the only case that is not a sublocus of the space \mathcal{S} . The equation of the curve from Table 3 of [20] is

$$y^2 = x(x^2 - 1)(x^4 + ax^2 + b)$$

In this case we have $J_3 = J_5 = J_7 = 0$. The defining equations of this space are two polynomials in J_2, J_4, J_6, J_8 . We make them part of the "genus3" package in [29]. It is worth noting that both a and b can be expressed as rational functions in t_1, \dots, t_6 . Hence, in this case the field of moduli is a field of definition.

5.3. 1-dimensional strata.

5.3.1. *The automorphism group is isomorphic to $\mathbb{Z}_2 \times D_8$:* Given a curve C in the $\mathbb{Z}_2 \times D_8$ locus, from Table 1, it has equation:

$$(27) \quad Y^2 = X^8 + aX^4 + 1,$$

where $a \neq \pm 2$. We calculate the invariants, t_1, t_2, \dots, t_6 and denote $t := a^2$. Then we have

$$\begin{aligned} t_1 &= 2 \frac{t(3t+980)^2}{(140+t)^3}, & t_2 &= 16 \frac{(t-196)^2}{(140+t)^2}, & t_3 &= 8 \frac{(t-196)^2}{(140+t)(3t+980)}, \\ t_4 &= 4 \frac{-196+t}{140+t}, & t_5 &= 4 \frac{t-196}{140+t}, & t_6 &= 128 \frac{(9t+980)(t-196)^3}{(140+t)^4} \end{aligned}$$

These invariants are not defined for $t = -140$ and $t = -\frac{980}{3}$. We can rewrite the above equations as

$$t = -28 \frac{5t_4 + 28}{t_4 - 4}$$

and the equations of this submoduli space are given by

$$t_1 = -\frac{175}{288}t_4^2 + \frac{125}{3456}t_4^3 + \frac{686}{27}, t_2 = t_4^2, t_3 = -6 \frac{t_4^2}{(5t_4 - 56)}, t_5 = t_4, t_6 = \frac{49}{3}t_4^3 + \frac{5}{12}t_4^4$$

Notice that we can get the equations in terms of J_2, \dots, J_8 very easily by substituting t_1, \dots, t_6 . Such equations would be valid in even in the cases when some of J_i are zero.

The equation of the curve can be written in terms of t as follows. Let $X \rightarrow a^{\frac{1}{4}}X$. Then the equation of the curve becomes

$$Y^2 = tX^8 + tX^4 + 1$$

Hence, for this family of curves the field of moduli is a field of definition.

Lemma 7. *Every genus 3 hyperelliptic curve with full automorphism group isomorphic to $Z_2 \times D_8$ has equation*

$$(28) \quad Y^2 = tX^8 + tX^4 + 1$$

for some $t \neq 0, 4$. If $t \neq -140, -\frac{980}{3}$ then

$$t = -28 \frac{5t_4 + 28}{t_4 - 4}$$

in terms of the absolute invariants and therefore the field of moduli is a field of definition.

5.3.2. *The automorphism group is isomorphic to D_{12} :* The equations of the curve is:

$$Y^2 = X(X^6 + aX^3 + 1)$$

We perform the following coordinate change $X \rightarrow a^{\frac{1}{3}}X$ and the equation of the curve becomes

$$Y^2 = X(tX^6 + tX^3 + 1),$$

where $t = a^2$. Then, we have

$$\begin{aligned} t_1 &= 9 \frac{t(4t + 245)^2}{(-35 + 2t)^3}, & t_2 &= \frac{(8t + 49)^2}{(-35 + 2t)^2}, & t_3 &= \frac{1}{3} \frac{(8t + 49)^2}{(-35 + 2t)(4t + 245)} \\ t_4 &= \frac{8t + 49}{-35 + 2t}, & t_5 &= \frac{8t + 49}{-35 + 2t}, & t_6 &= 3 \frac{(12t + 245)(8t + 49)^3}{(-35 + 2t)^4} \end{aligned}$$

We can eliminate t

$$t = \frac{7}{2} \frac{5t_4 + 7}{t_4 - 4},$$

and the equations for the submoduli space become

$$(29) \quad t_1 = \frac{686}{27} + \frac{125}{54}t_4^3 - \frac{175}{18}t_4^2, t_2 = t_4^2, t_3 = \frac{t_4^2}{5t_4 - 14}, t_5 = t_4, t_6 = \frac{65}{9}t_4^4 - \frac{98}{9}t_4^3$$

Lemma 8. *Every genus 3 hyperelliptic curve with full automorphism group isomorphic to D_{12} has equation*

$$(30) \quad Y^2 = X(tX^6 + tX^3 + 1)$$

for $t \neq 0, 4$. If $t \neq -\frac{35}{2}, -\frac{245}{4}$ then

$$t = \frac{7}{2} \frac{5t_4 + 7}{t_4 - 4}$$

in terms of the absolute invariants and therefore the field of moduli is a field of definition.

5.3.3. *The automorphism group is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_4$:* The equations of the curve is:

$$y^2 = (x^4 - 1)(x^4 + ax^2 + 1)$$

By a transformation $X \rightarrow a^{\frac{1}{2}}X$ the equation of the curve becomes

$$Y^2 = (tX^4 - 1)(tX^4 + tX^2 + 1)$$

Since this curve has an element of order 4 and therefore a factor of $X^4 - 1$ then $J_3 = J_5 = J_7 = 0$. In this case the absolute invariants t_1, \dots, t_6 are not defined. Hence we use the invariants i_1, \dots, i_5 as in Eq.(13). We have $i_1, i_3, i_5 = 0$ and

$$\begin{aligned} i_2 &= \frac{64}{25} \frac{t^2 + 9604 - 49t}{(28+t)^2}, \\ i_4 &= \frac{512}{125} \frac{(t-98)(t^2 - 637t + 9604)}{(28+t)^3}, \\ i_6 &= -\frac{512}{125} \frac{11t^4 - 12397t^3 + 1296540t^2 + 368947264 - 43294832t}{(28+t)^4} \end{aligned}$$

Hence, we get

$$t = 28 \frac{15625 i_2 i_4 - 152500 i_4 + 24375 i_2^2 + 1215200 i_2 - 2809856}{-15625 i_2 i_4 + 2500 i_4 + 245625 i_2^2 - 725600 i_2 + 401408}$$

and

$$(31) \quad \begin{cases} -81462500 i_4 + 927746400 i_2 - 963780608 - 256055625 i_2^2 - 1953125 i_4^2 \\ + 36093750 i_2 i_4 + 15187500 i_2^3 = 0 \\ -22689450000 i_6 - 4593393436800 i_2 + 4628074479616 + 52734375 i_6^2 + 8912109375 i_2^4 \\ + 1371093750 i_2^2 i_6 + 5788125000 i_2 i_6 + 1572126780000 i_2^2 - 215275375000 i_2^3 = 0 \end{cases}$$

5.4. **0-dimensional strata.** We first briefly go over the 0-dimensional cases.

5.4.1. *Case : $G \cong \mathbb{Z}_2 \times S_4$:* The equation of the curve is $y^2 = x^8 + 14x^4 + 1$ and its absolute invariants are

$$(t_1, t_2, t_3, t_4, t_5, t_6) = \left(\frac{15435}{8}, \frac{784}{25}, \frac{56}{25}, \frac{-28}{5}, \frac{28}{5}, \frac{7760032}{125} \right)$$

The next two cases correspond to curves with $J_3 = J_5 = J_7 = 0$. In both cases we use invariants i_2, i_4, i_6 as in Eq. (13).

5.4.2. *Case* : $G \cong U_6$: The equations of the curve is given by $y^2 = x(x^6 - 1)$ and its absolute invariants are $i_1 = i_3 = i_5 = 0$ and

$$i_2 = \frac{49}{25}, \quad i_4 = -\frac{343}{125}, \quad i_6 = \frac{7203}{125}$$

5.4.3. *Case* : $G \cong V_8$: The equations of the curve is $y^2 = x^8 - 1$ and its absolute invariants are $i_1 = i_3 = i_5 = 0$ and

$$i_2 = \frac{784}{25}, \quad i_4 = -\frac{21952}{125}, \quad i_6 = -\frac{307328}{125}$$

The following theorem determines relations among $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ for each group G such that $V_4 \hookrightarrow G$.

Theorem 4. *Let \mathcal{X} be a curve in $\mathcal{S} = \mathcal{M}_3^b \cap \mathcal{H}_3$. Then, one of the following occurs:*

- i) $\text{Aut}(\mathcal{X}) \cong \mathbb{Z}_2^3$ if and only if $\mathfrak{s}_4 - 2\mathfrak{s}_2^2 = 0$*
- ii) $\text{Aut}(\mathcal{X}) \cong \mathbb{Z}_2 \times D_8$ if and only if $\mathfrak{s}_2 = \mathfrak{s}_4 = 0$*
- iii) $\text{Aut}(\mathcal{X}) \cong \mathbb{Z}_2 \times \mathbb{Z}_4$ if and only if $\mathfrak{s}_4 + 2\mathfrak{s}_2^2 = 0$ and $\mathfrak{s}_3 = 0$.*
- iv) $\text{Aut}(\mathcal{X}) \cong D_{12}$ if and only if*

$$(32) \quad \begin{aligned} \mathfrak{s}_3 &= \frac{1}{75} (9\mathfrak{s}_2 - 224)(\mathfrak{s}_2 - 196) \\ \mathfrak{s}_4 &= -\frac{9}{125}\mathfrak{s}_2^3 + \frac{1962}{125}\mathfrak{s}_2^2 - \frac{840448}{1125}\mathfrak{s}_2 + \frac{9834496}{1125} \end{aligned}$$

Proof. Part i) and ii) are immediate consequences of the previous discussions. For part iii), we start with the curve \mathcal{X} with equation

$$Y^2 = (X^4 - 1)(X^4 + aX^2 + 1).$$

Transforming $X \rightarrow \epsilon_{16}X$ we have

$$Y^2 = X^8 - \epsilon_{16}^6 aX^6 + \epsilon_{16}^2 aX^2 + 1,$$

where ϵ_{16} is the 16-th root of unity. The dihedral invariants are

$$\mathfrak{s}_2 = a^2, \quad \mathfrak{s}_3 = 0, \quad \mathfrak{s}_4 = 2a^4$$

By eliminating a^2 we have that

$$\mathfrak{s}_4 + 2\mathfrak{s}_2^2 = 0, \quad \text{and} \quad \mathfrak{s}_3 = 0$$

Conversely, if the above equations hold then $a^2 + c^2 = 0$. Take a curve with equation as in Eq. (4) and compute i_2, i_4, i_6 . These invariants satisfy Eqs. (30). Hence, the curve is in the $(\mathbb{Z}_2 \times \mathbb{Z}_4)$ -locus.

For case iv), let \mathcal{X} be a curve with equation $Y^2 = X(X^6 + aX^3 + 1)$, where $a \neq 0, \pm 2$. By a transformation $X \rightarrow \frac{X+1}{X-1}$, \mathcal{X} has equation

$$Y^2 = X^8 + (5 - 9\lambda)X^6 + 3(\lambda + 1)X^4 + (5\lambda - 9)X^2 + \lambda$$

where $\lambda = \frac{a-2}{a+2}$, $\lambda \neq 0, \pm 1$. Then, by another transformation $X \rightarrow \sqrt[8]{\lambda}X$, we get the following curve

$$Y^2 = X^8 + \frac{(5 - 9\lambda)}{\lambda^{\frac{1}{4}}}X^6 + 3\frac{(\lambda + 1)}{\lambda^{\frac{1}{2}}}X^4 + \frac{(5\lambda - 9)}{\lambda^{\frac{3}{4}}}X^2 + 1$$

Computing the dihedral invariants:

$$\begin{aligned}\mathfrak{s}_2 &= \frac{1}{\lambda} (5 - 9\lambda)(5\lambda - 9) \\ \mathfrak{s}_3 &= \frac{3}{\lambda} (\lambda + 1) \left[(5 - 9\lambda)^2 + \frac{1}{\lambda} (5\lambda - 9)^2 \right] \\ \mathfrak{s}_4 &= \frac{1}{\lambda} (5 - 9\lambda)^4 + \frac{1}{\lambda^3} (5\lambda - 9)^4\end{aligned}$$

Eliminating λ , we get $\lambda = \frac{45}{106 - \mathfrak{s}_2}$ and

$$(33) \quad \begin{aligned}\mathfrak{s}_3 &= \frac{1}{75} (9 \mathfrak{s}_2 - 224) (\mathfrak{s}_2 - 196) \\ \mathfrak{s}_4 &= -\frac{9}{125} \mathfrak{s}_2^3 + \frac{1962}{125} \mathfrak{s}_2^2 - \frac{840448}{1125} \mathfrak{s}_2 + \frac{9834496}{1125}\end{aligned}$$

Conversely, let us assume that the equations in (33) hold. From the expressions of t_1, \dots, t_6 in Eqs. (18) and equations in (33) we eliminate $\mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4$ to get the equations of the D_{12} -locus in Eq. (29). The proof is complete. \square

REFERENCES

- [1] R. Alagna, *Le relazioni fra gl'invarianti d'una forma qualunque d'ottavo ordine*, Rendiconti del Circolo Matematico di Palermo **6** (1892), no. 1, 77–99 (Italian).
- [2] Jannis A. Antoniadis and Aristides Kontogeorgis, *On cyclic covers of the projective line*, Manuscripta Math. **121** (2006), no. 1, 105–130, DOI 10.1007/s00229-006-0028-4.
- [3] F. Bardelli and A. Del Centina, *Bielliptic curves of genus three: canonical models and moduli space*, Indag. Math. (N.S.) **10** (1999), no. 2, 183–190, DOI 10.1016/S0019-3577(99)80015-9. MR1816214 (2001m:14039)
- [4] Lubjana Beshaj, Valmira Hoxha, and Tony Shaska, *On superelliptic curves of level n and their quotients*, Albanian J. Math. **5** (2011), no. 3, 115–137.
- [5] Thomas Breuer, *Characters and automorphism groups of compact Riemann surfaces*, London Mathematical Society Lecture Note Series, vol. 280, Cambridge University Press, Cambridge, 2000. MR1796706 (2002i:14034)
- [6] Heinrich Burkhardt, *Untersuchungen aus dem Gebiete der hyperelliptischen Modulfunctionen*, Math. Ann. **38** (1891), no. 2, 161–224, DOI 10.1007/BF01199251 (German). MR1510670
- [7] ———, *Untersuchungen aus dem Gebiete der hyperelliptischen Modulfunctionen*, Math. Ann. **36** (1890), no. 3, 371–434, DOI 10.1007/BF01206368 (German). MR1510630
- [8] Maurizio Cornalba, *On the locus of curves with automorphisms*, Ann. Mat. Pura Appl. (4) **149** (1987), 135–151, DOI 10.1007/BF01773930 (English, with Italian summary). MR932781 (89b:14038)
- [9] Enrico Arbarello, Maurizio Cornalba, and Phillip A. Griffiths, *Geometry of algebraic curves. Volume II*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 268, Springer, Heidelberg, 2011. With a contribution by Joseph Daniel Harris. MR2807457 (2012e:14059)
- [10] Igor V. Dolgachev, *Classical algebraic geometry*, Cambridge University Press, Cambridge, 2012. A modern view. MR2964027
- [11] I. Dolgachev and T. Shaska, *Bielliptic curves of genus 3*, work in progress.
- [12] Y. Fuertes and G. González-Diez, *Fields of moduli and definition of hyperelliptic covers*, Arch. Math. (Basel) **86** (2006), no. 5, 398–408, DOI 10.1007/s00013-005-1433-8. MR2229355 (2007f:14028)
- [13] P. Gaudry and É. Schost, *On the invariants of the quotients of the Jacobian of a curve of genus 2*, (Melbourne, 2001), Lecture Notes in Comput. Sci. vol. 2227, Springer, Berlin, 2001, pp. 373–386.
- [14] W. D. Geyer, *Invarianten binärer Formen*, Classification of algebraic varieties and compact complex manifolds, Springer, Berlin, 1974, pp. 36–69. Lecture Notes in Math., Vol. 412 (German).

- [15] J. Gutierrez, D. Sevilla, and T. Shaska, *Hyperelliptic curves of genus 3 with prescribed automorphism group*, Computational aspects of algebraic curves, Lecture Notes Ser. Comput. vol. 13, World Sci. Publ., Hackensack, NJ, 2005, pp. 109–123.
- [16] J. Gutierrez and T. Shaska, *Hyperelliptic curves with extra involutions*, LMS J. Comput. Math. **8** (2005), 102–115.
- [17] Bonnie Huggins, *Fields of moduli of hyperelliptic curves*, Math. Res. Lett. **14** (2007), no. 2, 249–262. MR2318623 (2009a:14040)
- [18] Reynald Lercier and Christophe Ritzenthaler, *Hyperelliptic curves and their invariants: geometric, arithmetic and algorithmic aspects*, J. Algebra **372** (2012), 595–636.
- [19] Reynald Lercier, Christophe Ritzenthaler, and Jeroen Sijsling, *Fast isomorphism of hyperelliptic curves and explicit descent*, arXiv:1203.5440.
- [20] K. Magaard, T. Shaska, S. Shpectorov, and H. Völklein, *The locus of curves with prescribed automorphism group*, Sürikaiseikikenkyūsho Kōkyūroku **1267** (2002), 112–141. Communications in arithmetic fundamental groups (Kyoto, 1999/2001).
- [21] T. Shaska, C. Shor, F. Thompson, and E Zhupa, *Computational aspects superelliptic curves* (work in progress).
- [22] T. Shaska, *Subvarieties of the hyperelliptic moduli determined by group actions*, Serdica Math. J. **32** (2006), no. 4, 355–374.
- [23] ———, *Computational aspects of hyperelliptic curves*, Computer mathematics, Lecture Notes Ser. Comput. vol. 10, World Sci. Publ., River Edge, NJ, 2003, pp. 248–257.
- [24] ———, *Determining the automorphism group of a hyperelliptic curve*, Proceedings of the 2003 International Symposium on Symbolic and Algebraic Computation, ACM, New York, 2003, pp. 248–254 (electronic).
- [25] ———, *Genus 2 curves with $(3, 3)$ -split Jacobian and large automorphism group*, Algorithmic number theory (Sydney, 2002), Lecture Notes in Comput. Sci. vol. 2369, Springer, Berlin, 2002, pp. 205–218.
- [26] ———, *Genus two curves covering elliptic curves: a computational approach*, Computational aspects of algebraic curves, Lecture Notes Ser. Comput. vol. 13, World Sci. Publ., Hackensack, NJ, 2005, pp. 206–231.
- [27] ———, *Some remarks on the hyperelliptic moduli of genus 3*, Comm. Algebra, to appear.
- [28] ———, *Genus 3 hyperelliptic curves with $(2, 4, 4)$ decomposable Jacobians*, work in progress.
- [29] ———, *Algebraic Curves* (2012), www.risat.org/algcurves.
- [30] T. Shaska and H. Völklein, *Elliptic subfields and automorphisms of genus 2 function fields*, Algebra, arithmetic and geometry with applications (West Lafayette, IN, 2000), Springer, Berlin, 2004, pp. 703–723.
- [31] Tetsuji Shioda, *On the graded ring of invariants of binary octavics*, Amer. J. Math. **89** (1967), 1022–1046.
- [32] Von Gall, *Ueber das vollständige System einer binären Form achter Ordnung*, Math. Ann. **17** (1880), no. 1, 139–152 (German).